

## 2004

Novel Cationic Ring-Opening Polymerization of Cyclodextrin: A Uniform Macrocyclic Monomer with Unique Character

M. Suzuki\*, O. Numata, T. Shimazaki

Macromol. Symp. 2004, 215, 255-265.

Three-Component, Negative-Type, Alkaline-Developable, Thermally Stable, and Photosensitive Polymer Based on Poly(2,6-dihydroxy-1,5-naphthalene), a Crosslinker, and a Photoacid Generator

K. Tsuchiya, Y. Shibasaki, M. Suzuki, and M. Ueda\*

J. Polym. Sci., Part A, Polym. Chem. 2004, 42, 2235-2240.

Tropone-Containing p-Conjugated Polymers: Annulation of Tropone onto Benzene Ring in the Conjugated Polymer

N. Nishioka, K. Takagi,\* T. Kinoshita, H. Kunisada, and Y. Yuki

J. Polym. Sci. Part A: Polym. Chem. 2004, 42(5), 1208-1215.

Coordination of Transition Metal Dithiolate Complex to 2,2'-Bipyridine-Based p-Conjugated Polymer

S. Kato, K. Takagi\*, K. Suzuki, T. Kinoshita, and Y. Yuki

J. Polym. Sci. Part A: Polym. Chem. 2004, 42(11), 2631-2639.

Tropone-Containing Conjugated Polymer, Part V Influence of Fused Benzene Ring upon Photophysical Properties

K. Takagi\*, K. Mori, H. Kunisada, and Y. Yuki

Polym. Bull. 2004, 52, 125-132.

## 2003

Solvent-Free Polycondensation of 1,3-Bis(hydroxyphenylmethyl)benzene. II. The Dramatic Difference in the Polymerization Behavior between Meso and Racemic Isomers

M. Suzuki\*, T. Sasaki, G. Matsuura, and M. Kimura

J. Polym. Sci., Part A, Polym. Chem. (Special Issue: Japan–US Seminar on Advanced Polymer Chemistry for the 21st Century) 2003, 41, 3564–3571.

Novel Solid–State Polymerization of Crystalline Monomer. Dehydrative Polycondensation of 1,3–Bis(hydroxyphenylmethyl)benzene

T. Sasaki, K. Hayashibara, and M. Suzuki\*

Macromolecules, 2003, 36, 279–281, [Erratum]; Macromolecules 2004, 37, 4332.

Preparation of a Unique Glucan with Large Intervals in Molecular Weight Distribution. Controlled Ring–Opening Polymerization of O–Permethylcyclodextrin

M. Suzuki\* and T. Shimazaki

Org. Biomol. Chem. 2003, 1, 604–608.

A New Negative–Type Photosensitive Polymer Based on Poly(2,6–dihydroxy–1,5–naphthalene), a Cross–Linker, and a Photoacid Generator

K. Tsuchiya, Y. Shibasaki, M. Suzuki, and M. Ueda\*

J. Photopolym. Sci. Tech. 2003, 16, 285–286.

Acid–Doping to Tropone–Containing Conjugated Polymer

K. Takagi,\* K. Mori, T. Kinoshita, H. Kunisada, and Y. Yuki

Chem. Lett. 2003, 32(6), 552–553.

Synthesis of Star Polystyrene by Radical Polymerization with 1,2,4,5–Tetrakis(p–tert–butylphenylselenomethyl)benzene as a Novel Photoiniferter

T. S. Kwon,\* K. Takagi, H. Kunisada, and Y. Yuki

Eur. Polym. J. 2003, 39 (7), 1437–1441.

Allylnickel–Catalyzed Living Coordination Polymerization of Allenes with Ester Functionality

K. Takagi, I. Tomita,\* and T. Endo

Polym. Bull. 2003, 50(5–6), 335–342.

2002

Facile Preparation of Mono-2-O-Modified  
Eicosa-O-methylcyclomaltoheptaoses ( $\beta$ -cyclodextrins)

M. Suzuki\* and Y. Nozoe

Carbohydrate Res. 2002, 337, 2393–2397.

New Positive-type Photosensitive Polymer Based on  
Poly(2,6-dihydroxy-1,5-naphthylene) and Diazonaphthoquinone

Y. Sasada, Y. Shibasaki, M. Suzuki, and M. Ueda\*

J. Polym. Sci., Part A, Polym. Chem. 2002, 40, 393–398.

Convenient Synthesis of Poly(2,6-dihydroxy-1,5-naphthylene) by  
Cu(II)-amine Catalyzed Oxidative Coupling Polymerization

Y. Sasada, Y. Shibasaki, M. Suzuki, and M. Ueda\*

Polymer 2002, 44, 355–360.

Novel Solid-State Polycondensation I. Oxidative-Coupling Polymerization of  
2,6-Dihydroxynaphthalene

M. Suzuki\* and Y. Yatsugi

Chem. Commun. 2002, 162–163.

$\pi$ -Conjugated Oligomers Containing Tropone in the Main Chain: Synthesis,  
Characterization, and Optical Properties

K. Takagi,\* Y. Nishikawa, N. Nishioka, H. Kunisada, and Y. Yuki

J. Polym. Sci. Part A: Polym. Chem. 2002, 40(22), 3927–3937.

Synthesis of Diblock Copolymers by End Functional Polystyrene Containing  
Phenylseleno Groups as Polymeric Photoiniferter

T. S. Kwon,\* K. Takagi, H. Kunisada, and Y. Yuki

J. Macromol. Sci., Pure and Appl. Chem. 2002, A39(9), 991–1006.

The Adsorption of Guest Molecule onto Self-Assembled Monolayer of  
 $\alpha$ -Helical Polypeptide

I. Aoki, N. Nishioka, K. Takagi, Y. Yuki, and T. Kinoshita\*

II. Trans. of Mater. Res. Soc. Jpn. 2002, 27(3), 559–562.

## 2001

Cyclodextrin as Macrocyclic Monomer. Cationic Ring–Opening  
Polymerization of O–Permethylcyclodextrins

M. Suzuki\*, O. Numata, and T. Shimazaki

Macromol. Rapid Commun. 2001, 22, 1358–1361.

Synthesis and Polymerization of Isopropenyl–1,3,5–triazines Carrying a Long  
Alkyl Chain with Oxyethylene Spacer and the Side Chains Crystallization of  
Obtained Comb–like Polymers

H. Kunisada,\* T. Ito, K. Takagi, and Y. Yuki

J. Macromol. Sci., Pure and Appl. Chem. 2001, A38(4), 417–427.

Radical Polymerization of Methyl Methacrylate with Diphenyl Diselenide  
under Thermal or Photoirradiational Conditions

T. S. Kwon, K. Suzuki, K. Takagi, H. Kunisada, and Y. Yuki\*

J. Macromol. Sci., Pure and Appl. Chem. 2001, A38(5–6), 591–604.

Synthesis of ABA Type Triblock Copolymers by Radical Polymerization with  
1,4–Bis(p–tert–butylphenylselenomethyl)benzene as a Photoiniferter

T. S. Kwon, K. Takagi, H. Kunisada, and Y. Yuki\*

J. Macromol. Sci., Pure and Appl. Chem. 2001, A38(5–6), 605–626.

Synthesis of PPV–Based Conjugated Oligomer Containing Tropone in Main  
Chain

K. Takagi,\* Y. Nishikawa, H. Kunisada, and Y. Yuki

Chem. Lett. 2001, (12), 1244–1245.

## 2000

Hydrogen–transfer polymerization behavior of N–acrylamide

T. Iwamura, I. Tomita, M. Suzuki, and T. Endo\*

J. Polym. Sci., Part A, Polym. Chem. 2000, 38 (3), 430–435.

Reaction of p-toluenesulfonyl isocyanate with polymers having amide moieties and

hydrolysis of the obtained polymer

T. Iwamura, I. Tomita, M. Suzuki, and T. Endo\*

J. Polym. Sci., Part A, Polym. Chem. 2000, 38 (18), 3440–3449.

Ni-Catalyzed Living Polymerization of Allenes; Precision Polymers and Nano-Organized Systems

M. Suzuki, I. Tomita, and T. Endo\*, Ed. by T. Kunitake, S. Nakahama, S. Takahashi, and N. Toshima, Kodansha Ltd.: Tokyo, 2000, 19–22

Synthesis of Triazine Dendrimers

K. Takagi,\* K. Uchikura, T. Hattori, H. Kunisada, and Y. Yuki

Kobunshi Ronbunshu 2000, 57(10), 646–651.

Synthesis of Macromonomer Using End Functional Polystyrene Prepared from p-Methoxybenzyl p-Trimethylsilylphenyl Selenide as Photoiniferter

T. S. Kwon, K. Takagi, H. Kunisada, and Y. Yuki\*

J. Macromol. Sci., Pure and Appl. Chem. 2000, A37(11), 1461–1473.

Synthesis of Branched Polystyrene by Photopolymerization of Selenium-Containing Styrene Monomer

K. Takagi,\* Y. Nishikawa, T. S. Kwon, H. Kunisada, and Y. Yuki

Polym. J. 2000, 32(11), 970–973.

Triazine Dendrimers by Divergent and Convergent Methods

K. Takagi,\* T. Hattori, H. Kunisada, and Y. Yuki

J. Polym. Sci. Part A: Polym. Chem. 2000, 38(24), 4385–4395.